

EFFECT OF DIET DILUTION IN THE STARTER PERIOD ON PERFORMANCE AND CARCASS CHARACTERISTICS OF BROILER CHICKS

M. REZAEI*¹, A. TEIMOURI², J. POURREZA³, H. SAYYAHZADEH¹ and P. W. WALDROUP⁴

1- Dept. of Anim. Sci, Faculty of Agriculture, Univ. of Mazandaran, P. O. Box 578, Sari, Iran.

2-Former MSc. student, Faculty of Agriculture, Univ. of Mazandaran, P. O. Box 578, Sari, Iran

3- Dept. of Anim. Sci, Faculty of Agricultur, Univ. of Tech Isfahan, P. O. Box 8415683111, Isfahan, Iran.

4- Dept. of Poult. Sci, Univ. of Arkansas, Fayetteville, AR 72701, USA.

E-mail::m.rezaei@umz.ac.ir

Manuscript received: November 7, 2005; Reviewed: May 13, 2006; Accepted for publication: June 9, 2006

ABSTRACT

The effect of energy and protein dilution in the starter period (8 to 14 days) of age, on performance and carcass characteristics of 360 Arian male chicks was studied in a completely randomised design. This experiment consisted of 6 treatments, 4 replicates, with 15 chicks per replicate. In order to dilute the diets six levels (0, 4, 8, 12, 16, and 20) percent of ground wood charcoal was used. Chicks were fed with starter and grower diet from 15 to 21 and 22 to 42 days of age respectively. During the experiment feed intake, body weight gain, feed conversion ratio were measured weekly. Mortality was measured throughout the experiment. At 43 day of age 4 chicks each treatment was selected and carcass characteristics were measured. The results indicated that dilution of diet from 8 to 14 days of age increased feed intake in this period but the differences were not significant. With increasing dilution rate body weight gain of chicks significantly decreased in comparison to control group ($P < 0.05$). Due to compensatory growth after restricted period, there was not significant difference in body weight among restricted and control groups at 42 days of age. There was not significant difference among the treatments for feed conversion ratio in whole period of the experiment (8 to 42 day). Similarly, there was not significant difference among the treatments for caloric conversion ratio in whole period of the experiment (8 to 42 day). Mortality rate in restricted groups was significantly lower than control group ($P < 0.05$). Diet dilution hadn't significant effect on carcass, breast meat, drumsticks, thighs, liver, intestine, abdominal fat percentages. The results of the present study indicated that broiler chicks could withstand a 7-day period (from 8 to 14 days) feed restriction with ground wood charcoal in early age without loss in performance.

Keywords: broiler chicks, diet dilution, wood charcoal, performance

INTRODUCTION

Feeding strategy in growing broiler chickens should be to produce animals with maximum lean body mass, highest feed conversion ratio and maximum body weight. Continuous genetic selection and improvement in nutrition have led to a very fast growth rate in modern strains (11,12,17,18,19,29). To the extent that the time required growing a broiler chicken to 2 kg has decreased from 63 d to 37 d (34). Research shows that the improved growth rate results from a large increase in early postnatal growth rate (11). A number of problems, namely increased body fat deposition, high incidence of metabolic disorders, high mortality, and high incidence of skeletal diseases accompany the early-life growth rate. To tackle with these problems early nutrient restriction programmers were used (9,10,11,12,29). Urdaneta-Rincon and Leeson, (30), Saleh et al. (22) reported that feed restriction and diet dilution cause temporary reduced growth rate, assuming that normal weight is realized at market age, feed efficiency should be improved (9). Early feed restriction of broiler chickens is usually applied in order to induce catch-up growth and improved efficiency of feed utilization (28). Experimental results

show considerable variation in the responses of chicken to feed restriction (26, 29, 34). Reasons for the relative success or failure of feed-restricted chickens to achieve full body weight recovery following realimentation are still unknown (28). Dilution of the diet with oat hulls, rice bran, cellulose or inert filler can be a rather easy way to induce growth retardation (12). Leeson et al. (10) found a complete recovery of body weight at the age of 42 d, after diet dilution with ground rice hulls up to 55 % from 4 to 11 d of age. The overall efficiency of feed utilization was not affected. However, there was an indication of reduced abdominal fat content for males at 56 days of age. Lee and Leeson (9) reported that male broiler fed diets diluted up to 50 % with oat hulls from 7 to 14 days, caught up body weight by 8 wk of age but had abdominal fat levels similar to control birds. Breast meat yield also was decreased. As far as we are aware, wood charcoal has not been tested as inert filler in broiler chickens diets. In this experiment, wood charcoal was used as new inert filler. Therefore, the present study was conducted to evaluate the effect of feed restriction with using wood charcoal on performance, carcass characteristics and mortality of Arian (Hybro) male broiler chicks.

Table 1. Composition (%) and calculated nutrient content of diets

Ingredients and Composition (%)	Starter		Test diet				Grower
	1(control)	2	3	4	5	6	
Corn	54.40	52.16	49.94	47.56	45.07	42.97	64.15
Soybean meal	40.16	38.34	36.46	34.71	33.14	31.03	31.88
Wood charcoal	---	4.00	8.00	12.00	16.00	20.00	---
Soy oil	2.00	1.96	1.92	1.91	1.93	1.86	1
Dicalcium phosphate	1.30	1.33	1.37	1.41	1.44	1.49	0.84
Oyster shell	1.28	1.27	1.27	1.26	1.25	1.25	1.35
Salinomycin	0.05	0.05	0.05	0.05	0.05	0.05	0.05
L-Lysine	---	0.05	0.12	0.2	0.18	0.33	---
Salt	0.30	0.30	0.30	0.30	0.30	0.30	0.29
DL-Methionine	0.09	0.13	0.16	0.19	0.22	0.26	0.01
Mineral premix ¹	0.25	0.25	0.25	0.25	0.25	0.25	0.25
Vitamin permix ²	0.25	0.25	0.25	0.25	0.25	0.25	0.25
Total	100	100	100	100	100	100	100
Calculated analysis							
ME (KcaL /Kg)	2900	2784	2668	2552	2436	2320	2950
Crude protein	20.9	20.0	19.2	18.39	17.55	16.7	18.4
Calcium (%)	0.89	0.89	0.89	0.89	0.89	0.89	0.79
Available phosphorus (%)	0.40	0.40	0.40	0.40	0.40	0.40	0.30
Sodium (%)	0.14	0.14	0.14	0.14	0.14	0.14	0.14
Arginine (%)	1.58	1.58	1.44	1.37	1.31	1.23	1.36
Lysine (%)	1.30	1.28	1.28	1.28	1.21	1.26	1.08
Methionine (%)	0.45	0.47	0.49	0.50	0.52	0.54	0.34
Methionine+cystine (%)	0.81	0.82	0.82	0.82	0.82	0.82	0.66

¹Mineral mix supplied the following per kg of diet: Cu, 20 mg; Fe, 100 mg; Mn,100 mg; Se, 0.4; Zn, 169.4 mg.

²Vitamin mix supplied the following per kg of diet: vitamin A, 18000 IU;vitamin D₃, 4000 IU; vitamin E, 36mg; vitamin K₃,4 mg; vitamin B₁₂,0.03 mg; thiamine, 1.8 mg; riboflavin, 13.2 mg; pyriod oxin, 6 mg; niacin, 60 mg; calcium pantothenate, 20 mg; folic acid, 2 mg; biotin, 0.2;cholin chloride, 500 mg.

MATERIAL AND METHOD

In this experiment, diet dilution was achieved by substitution of ground wood charcoal for the major ingredients in the diet (Diets 2,3,4,5, and 6). The experimental diets, containing 0 (zero), 4, 8, 12, 16, 20 % ground wood charcoal were formulated to be isonitrogenous and isocaloric by using maize, soybean meal, soy oil as main ingredients (Table 1). Chicks were fed with a commercial starter diet from 1 to 7 days of age. At d 8 chicks were distributed among 24 experimental pens (1×2 m). Between 15 to 21 days of ages, all birds were fed the regular starter diet (Table 1). After this period, chicks were fed with a regular grower diet up to 42 days of age (Table 1). All diets were formulated to meet the nutrient requirements according to NRC (14). Ground wood charcoal contains 94.1 % DM, 14.5 % ash, 75.7 % CF, 2.16 % CP, and 1.08 % EE, and 0.66 % NFE. A total of 360 male broiler chicks (Arian) were obtained from a commercial hatchery. During the experiment weight gain, feed intake and feed conversion ratio were measured weekly. Mortality was measured throughout the experiment. At the end of the experiment (43 d) 1 bird from each pen with body weight close to the pen average selected for carcass analyses. After feed withheld for 9h, the selected birds were transported to the university pilot for processing. The chickens were slaughtered by cervical dislocation to determine the carcass characteristics.

Data of this experiment were analyzed by analysis of variance using General Linear Models (GLM) procedures of SAS (25). When necessary, the means were compared by Duncan's Multiple Range Test (4). Percentage data for processing effects were converted to Arcsine before analysis. Mortality data were transformed to $\sqrt{n+1}$ prior to analysis. The level of significance was reported at $P<0.05$.

RESULTS AND DISCUSSION

Effect of diluted diets on feed intake, body weight gain, feed conversion ratio, calorie conversion ratio, carcass characteristics and mortality are presented in Tables 2 and 3.

Feed intake

During the period of diet dilution, birds attempted to maintain nutrient intake by consuming more feed, thus increasing degrees of diet dilution resulted in graded increase in feed intake (Table 2). This is in contrast with the findings of Leeson et al., and Yussefi Kelaricolaei et al. (10, 31). If wood charcoal is excluded from the calculation of feed intake (assumed indigestible) the feed intake of birds in treatments 4, 5 and 6 reduced ($P<0.05$).

During the 15 to 42 day (realimentation period), diet dilution hadn't significant effect on this trait. Similarly feed intake between 8 to 42 days of age wasn't affected by the diet dilution in 8 to 14 days of age. Results of this experiment are in agreement with the finding of other studies (9, 31). It seems that due to using low levels of wood charcoal in diets, there weren't significant difference among treatments in whole period of the experiment for this trait.

Body weight gain

With increasing Diet dilution there was a corresponding reduction in body weight gain by 7% in treatment 6. When birds resumed eating the regular undiluted starter diet after 14 days of age, prior diet dilution hadn't significantly effect on BWG. There weren't significant difference among treatments in whole period of the experiment. Results of this study are in agreements with findings of Leeson et al. (10), Lee and Leeson (9), Yussefi Kelaricolaei et al., (31) Lippens et al. (12). With increasing feed restriction (treatment 6) birds could not obtain the nutrients that are necessary for the normal growth.

Feed conversion ratio

During the period of diet dilution (8 to 14 days of ages), no significant difference was observed between birds fed with diluted and control diets. If wood charcoal is again assumed to be indigestible, then adjusted feed conversion ration was improved in treatments 4, 5 and 6 ($P<0.05$). There wasn't significant difference among treatments for this trait in whole period of the experiment (8 to 42d). With excluding wood charcoal in diets feed conversion ratio was improved in treatment 2, 4 in whole period (8 to 42). This results in agreement with the findings of Lee and Lesson (9), Yussefi Kelaricolaei et al. (31). The improvement in feed conversion ratio observed in feed restricted chickens has been attributed to reduce over all maintenance requirements caused by a transient decrease in basal metabolic rate. However, the improved feed conversion ratio can also be related to higher feed and to the enlargement of the gastrointestinal tract that occurs after the restriction period, when the birds are fed ad libitum.

Calorie conversion ratio

This measurement may be more economically important estimate of the ability of the birds to utilize their diets that is feed conversion. During the period of diet dilution (8 to 14 days), calorie conversion ratio was improved in treatment 4 and 5 ($P<0.05$). If wood charcoal is again assumed to be indigestible, then adjusted calorie conversion ratio was significantly reduced for treatment 3, 4, 5 and 6. There wasn't significant different among

Table 2. Effect of diluted diets on feed intake, body weight gain, feed conversion ratio, calorie conversion ratio of male broiler chicks

treatment	1	2	3	4	5	6	SEM
Feed intake (g)							
8 to 14* d	349	364	345	350	360	358	12.85
8 to 14** d	349 ^a	350 ^a	317 ^{ab}	310 ^b	302 ^b	286 ^b	11.5
8 to 42* d	4497 ^{ab}	4372 ^b	4574 ^{ab}	4368 ^b	4630 ^a	4504 ^{ab}	64
8 to 42** d	4497 ^{abc}	4358 ^{bc}	4546 ^{ab}	4329 ^c	4572 ^a	4436 ^{abc}	64.5
15 to 42 d	4027	4021	4269	4010	4262	4370	88.5
Body weight gain (g)							
8 to 14 d	256 ^a	265 ^a	252 ^a	267 ^a	258 ^a	238 ^b	5.5
8 to 42 d	1901	1894	1921	1967	1930	2004	71
15 to 42 d	1639	1671	1679	1638	1710	1765	71.85
Feed conversion ratio							
8 to 14 *d	1.345	1.355	1.360	1.300	1.382	1.470	0.05
8 to 14** d	1.345 ^a	1.320 ^{ab}	1.250 ^{abc}	1.140 ^{bc}	1.125 ^{bc}	1.150 ^{bc}	0.05
8 to 42* d	2.227	2.307	2.385	2.250	2.467	2.200	0.11
8 to 42** d	2.227 ^a	2.142 ^b	2.295 ^a	2.120 ^b	2.250 ^{ab}	2.205 ^{ab}	0.04
15 to 42 d	2.422	2.292	2.525	2.440	2.475	2.385	0.04
Calorie conversion ratio (ME kcal/kg)							
8 to 14* d	3939 ^a	3838 ^{ab}	3639 ^{abc}	3332 ^c	3392 ^{bc}	3480 ^{abc}	149.12
8 to 14** d	3939 ^a	3687 ^{ab}	3346 ^{bc}	2959 ^{cd}	2848 ^d	2777 ^d	138.83
8 to 42* d	6529	6395	6524	6160	6461	6329	159
8 to 42** d	6529	6234	6618	6244	6380	6228	124
15 to 42 d	7219	7011	7524	7354	7160	7165	331.8

Means within rows with no common superscripts differ significantly ($P < 0.05$).

** Feed intake, Feed conversion ratio, Calorie conversion ratio excluded the wood charcoal

* Feed intake, Feed conversion ratio, Calorie conversion ratio do not excluded the wood charcoal

Treatment 1, 2, 3, 4, 5, 6 contain 0, 4, 8, 12, 16, and 20 percent of ground wood charcoal, respectively

Table 3. Effect of diluted diets on carcass characteristics and mortality of male broiler chicks

Carcass characteristics and Mortality	Treatment						SEM
	1	2	3	4	5	6	
Live weight (g)	1928	1826	1851	1823	1951	1915	9.2
Carcass weight (g)	1305	1249	1232	1294	1298	1318	8.48
Carcass (%LW ¹)	67.68	68.37	66.68	71.24	67.80	68.78	1.69
Breast (% CW ²)	25.43	26.05	26.74	23.47	26.08	25.64	1.12
Thighs (% CW)	14.60	14.56	14.25	14.85	14.98	14.24	0.67
Drumsticks (% CW)	13.09	13.18	14.02	14.10	13.68	13.58	0.69
Liver (% CW)	4.17	4.32	3.98	4.23	4.54	4.39	0.54
Intestine (% CW)	8.52	9.02	9.55	9.09	9.07	8.89	0.91
abdominal fat content (% CW)	2.79	2.60	2.80	2.67	2.38	2.60	0.65
Mortality (8 to 42 %)	8.278 ^a	0.00 ^b	3.503 ^b	3.503 ^b	3.225 ^{ab}	0.00 ^b	0.006

Means within rows with no common superscripts differ significantly ($P < 0.05$).

¹Live weight

²Carcass weight

Treatment 1, 2, 3, 4, 5, 6 contain 0, 4, 8, 12, 16, and 20 percent of ground wood charcoal, respectively.


treatments for this trait in whole period of the experiment (8 to 42). The results of the present study are in agreement with finding of other researchers (22, 23).

Mortality

As expected, mortality following diet dilution was significantly lower for bird that was restricted at 8 to 14 days of age. Many investigators have reported a reduction in mortality rate following feed restriction (1,20,22,23,29). This could provide the greatest economic incentive for implementing early feed restriction by allowing for more birds to be marketed from a flock. In present study, mortality was reduced in restricted groups of chicks. The two main metabolic disorders affecting contemporary broiler chickens are ascites and SDS. These pathologic conditions are closely related to rapid growth rate and increase feed intake. Feed restriction slows down fast growth to reduce late mortality, including preascites and ascites (22,23,27,29). Feed restriction decreased mortality caused by “sudden death syndrome” (11). In this experiment implementation of feed restriction significantly reduce the prevalence of these disorders, which is in agreement with observations of Julian (7), Gonzales (5) but in contrast with Robinson et al. (20) and McGovern (13) and Urdaneta-Rincon, and Leeson (30). Effect of diluted diets on survivability rate of male broiler chicks is shown in Figure 1.

Carcass characteristics

Carcass weight expressed as the percent of live weight was not affected by feed restriction. Thighs, Drumsticks, Breastmeat, Intestine, Liver, Abdominal fat pad percentage were not affected. One of the most controversial aspects of early feed restriction programs has been lack of a consistent effect on abdominal fat pad (11). A reduction in abdominal fat content with concomitant reduction in body weight were found by Plavink and Hurwitz (17,18,19) and Jones and Farrell (6). Other investigators have reported reductions in abdominal fat pad due to early life feed restriction but a small reduction in final body weight (3). Other researchers (9, 10,23,24,30) were not able to show a clear effect. Bean et al (2) reported that feed efficiency was improved, but the amount of abdominal fat in male broilers was significantly increased. Zubair and Leeson (35) showed that feed-restricted birds usually had a numerically smaller abdominal fat pad. This agrees with the current results. The fact that there was no significant reduction in abdominal fat deposition in this experiment suggests that even feed-restricted broiler chickens are still overeating and that the level of feed intake may control de novo lipogenesis (21). It seems that a more severe time of feed restriction is necessary to significantly reduced abdominal fat pad. Carcass weight at 43d was not affected by diet dilution. None



^{a,b} Means with no common letters differ significantly (P<0.05). Treatment 1, 2, 3, 4, 5, 6 contain 0, 4, 8, 12, 16, and 20 percent of ground wood charcoal, respectively.

Figure 1. Effect of diluted diets on survivability rate of male broiler chicks

of the other carcass parts were significantly influenced by the feed restriction. In contrast to the situation with carcass or abdominal fat content, there are few results on effects of feed restriction on carcass characteristics. Leeson et al (10), Zubair and Leeson (32) and Palo et al (15), Lippens et al. (11), Yussefi Kelaricolaei et al., (31) could not demonstrated any effect of feed restriction in dressing percentage. No effect was found on other carcass characteristics by Leeson et al. (10), Zubair and leeson (33), Palo et al (16), Lippens et al.(11), Yussefi Kelaricolaei et al. (31), and this confirmed by our own results. Other researches found no significant effect of feed restriction on the percentage of thighs or drumsticks (11,22). Intestine, Liver percentage were not affected by feed restriction. These results are consistent with Palo et al. (15) and are contrast with Zubair and Leeson (33). In conclusion, feed restriction as used in the current trail give some indication of being a practical tool to reduce losses due to metabolic disease without deteriorating performance and carcass characteristics. Evidently the broiler chicken can withstand a 7-days period of under nutrient without any loss in market weight. More researches however are needed to confirm this.

CONCLUSION

The use of up to 20 % wood charcoal for dilution diet in starter broiler diet from 8 to 14 days of age, could reach the same performance at 42 days of age as in case of using control diet (containing 0 % wood charcoal). In this case, mortality in chicks fed with diluted diets significantly reduced in comparison with control group. Use of wood charcoal in starter diets hadn't any adverse effect on carcass characteristics.

ACKNOWLEDGEMENT

The authors would like to express their appreciation to Faculty of Agriculture, Mazandaran University for their financial support. The assistance of A. Naghdi., B. Shoghi, several graduate students from Faculty of Agriculture and the staff of Faculty of Agriculture is gratefully acknowledged.

REFERENCES

- [1] Arce J., Berger M., Lopez Coello C., Control of ascites syndrome by feed restriction techniques. *J. Appl. Poult. Res.* (1992) 1: 1 – 5.
- [2] Beane W. L., Cherry J. A., Weaver JR W. D., Intermittent light and restricted feeding of broiler chickens. *Poult. Sci.* (1979). 58: 567 – 571.
- [3] Cabel M. C., Waldroup P. W., Effect of different nutrition-restricted programs early in life on broiler performance and abdominal fat content. (1990). *Poult. Sci.* 69: 652 – 66.
- [4] Duncan D. B., Multiple ranges and Multiple F-test *Biometrics.* (1955) 11: 1-42.
- [5] Gonzales E., Buyes J., Lodd M. M, Takita T. S., Buys N., Decuypere E., Performance, incidence of metabolic disturbances and endocrine variables of food-restricted male broiler chickens. *Br. Poult. Sci.*(1989a) 39: 671 – 678.
- [6] Jones G. P. D., Farrell D. J., Early-life food restriction of broiler chicken I. Methods of application, amino acid supplementation and the age at which restriction should commence. *Br. Poult. Sci.* (1992). 33: 579 – 587.
- [7] Julian R. J., Causes and prevention of ascites in broilers. *Zootec. Int.* (1997) 4: 52- 53.
- [8] Kutlu H. R, Unsal I., Gorgulu M., Effects of Providing dietary wood (oak) charcoal to broiler chicks and laying hens. *Anim. Feed. Sci.Tech.* (2001) 90:213 – 226.
- [9] Lee K. H., Leeson S., Performance of broilers fed limited quantities of feed or nutrient during seven to fourteen days of age. (2001) *Poult. Sci.* 80: 446 – 454.
- [10] Leeson S., Summers J. D., Caston L. J., Diet dilution and compensatory growth in broilers. *Poult. Sci.* (1991) 70: 867-873.
- [11] Lippens M., Room G., Groote G.De., Decupere E., Early and temporary quantitative food restriction of broiler chickens . 1. Effects performance characteristics, mortality and meat quality. *Br. Poult. Sci.* (2000) 41: 343 – 354.
- [12] Lippens M., Huyghebaert G., Tuyt O.V., Groote G.De., Early and temporary Qualitative, autonomous feed restriction of broiler chickens. Effect on performance Characteristics, mortality, carcass and meat quality. *Arch. Geflugelkd.* (2002) 67: 49 – 56.
- [13] McGovern R. H., Robinson F. E., Feddes and J. J. R., Advences in ascites research. *Zootc. Int.* (1997) 8: 16 - 17.
- [14] National Research Council., Nutrient Requirements of poultry 9th rev. edition. *Natl. Acad. Press.* Washington. D C, (1994).
- [15] Palo P. E., Sell J. L., Piquer, F. J., Vilaseca L., Soto-Salanova M. F., Effect of early nutrient restriction on broiler chicken.I. performance and development of the gastrointestinal tract. *Poult. Sci.* (1995a) 74: 88 – 101.

- [16] Palo P. E., Sell J. L., Piquer F. J., Vilaseca L., Soto-Salanova M. F., Effect of early nutrient restriction on broiler chicken. I. Performance and digestive enzyme activities. *Poult. Sci.* (1995b) 74: 1470 – 1483.
- [17] Plavnik I., Hurwitz S., The performance of broiler chicks during and following a severe feed restriction at an early age. *Poult. Sci.* (1985) 64: 348 - 355.
- [18] Plavnik I., Hurwitz S., Response of broiler chickens and turkey poults to food restriction of varied severity during early life. *Br. Poult. Sci.* (1991) 32: 343 – 352.
- [19] Plavnik I., Hurwitz S., Early feed restriction in male turkeys. Growth pattern, feed efficiency and body composition. *Poult. Sci.* (1988) 67: 1407- 1413.
- [20] Robinson F. E., Classen H. L., Hanson J. A., Onderka D. K., Growth performance, feed efficiency and the incidence of skeletal and metabolic disease in full – fed and feed-restricted broiler and roaster chickens. *J. Appl. Poult. Res.* (1992. 1) : 33 – 41.
- [21] Rosebrough R. W., McMurtry J. P., Energy repletion and lipid metabolism during compensatory gain in broiler chickens. *Growth Dev. Aging.* (1993). 57: 73 - 83
- [22] Saleh, K., Attia Y. A., Younis H., Effect of feed restriction and breed on compensatory growth, abdominal fat and some production traits of broiler chicks. *Archiv. Geflugelkd.* (1996) 64(4): 153- 15.
- [23] Saleh E. A., Watkins S. E., Waldroup A. L., Waldroup P. W., Comparison of Energy feeding programs and early feed restriction on live performance and carcass quality of large male broiler grown for further processing at 9 to 12 weeks of age. *Int. J. Poult. Sci.* (2004) 3: 61 - 69.
- [24] Saleh E. A., Watkins S. E., Waldroup A. L., Waldroup P. W., Effects of early quantitative feed restriction on live performance and carcass composition of male broilers grown for Further processing. *J. Appl. Poult. Res.* (2005) 14: 87 – 93.
- [25] SAS Institute. SAS User'Guide: Statistic. Edition. SAS Institute Inc, Carry, NC. USA. (2001)
- [26] Scheideler S. E., Baughman G. R., Computerized early feed restriction Programs for various strains. *Poult. Sci.* (1993) 72: 236 – 242.
- [27] Summers J. D., Sparrt D., Atkinson J. L., Restricted feeding and compensatory growth for broiler. *Poult. Sci.* (1990) 69: 1855 - 1861
- [28] Susblia, J. P., Tarvid I., Gow C. B., Frankel T. L., Quantitative feed restriction or Meal-feeding of broiler chicks alters functional development of enzymes for protein Digestion. *Br. Poult. Sci.* (2003) 14: 698 - 709.
- [29] Tumova E., Skirvan M., Skirvanova V., Kacerovska L., Effect of early feed restriction on growth in broiler chickens, turkeys and rabbits. *Czech J. Anim. Sci.* (2002) 47: 418-428.
- [30] Urdaneta-Rincon M., Leeson S., Quantitative and qualitative feed restriction on growth Characteristics of male broiler chickens. *Poult. Sci.* (2002) 81: 679 – 688.
- [31] Yussefi Kelaricolaii, K., Rezaei M., Kamyab A., Performance of broiler chickens during and following feed restriction at an early age. *Br. Poult. Sci.* (2001) 42(suppl.1).111(abst).
- [32] Zubair A. K., Leeson S., Effect of early feed restriction and realimentation on heat production and changes in size of digestive organs of male broilers. *Poult. Sci.* (1994a) 73: 529 - 538.
- [33] Zubair A. K., Leeson S., Effect of varying period of early nutrition restriction growth compensation and carcass characteristics of male broilers. *Poult. Sci.* (1994b) 73: 129 - 136.
- [34] Zubair A. K., Leeson S., Compensatory growth in the broiler chicken: A Review. *World's. Poult. Sci. J.* (1996a) 52: 189- 201.
- [35] Zubair A. K., Leeson S., Changes in body composition and adipocyte cellularity of male broilers subjected to varying degrees of early – life feed restriction. *Poult. Sci.* (1996b) 75: 719 – 728.

